

Zapain



Zapain is a compound painkiller that contains the active ingredients paracetamol and codeine. Paracetamol is known for its fever-reducing and pain-relieving qualities. Codeine is a suppressant which blocks pain signals and the cough reflex.

Prices may vary slightly when paying.

Zapain 30/500 mg 100 tabl. **EUR46** [Buy now](#)

Zapain 30/500 mg 100 caps. **EUR46** [Buy now](#)

Description

Product information What is Zapain? When is it used? How to use Zapain Dosage Side effects
Pregnancy/alcohol/driving ability

What is Zapain?

Zapain is a compound painkiller that contains the active ingredients paracetamol and codeine. Paracetamol is known for its fever-reducing and pain-relieving qualities. Codeine is a suppressant which blocks pain signals and the cough reflex. This combination makes Zapain a very powerful painkiller that works quickly and effectively.

When is it used?

Zapain can be used when paracetamol alone does not suffice to control the pain. It is prescribed to treat moderate pain symptoms, either with or without fever. Zapain is used to treat headache, pain and fever due to the flu, dry cough, toothache, rheumatic pain, post-operative pain, pain from injuries or pain caused by other factors.

How to use Zapain

Zapain is available in tablet or capsule form and should be swallowed whole, without chewing, with some water. It is best to take Zapain during meals to avoid stomach distress. Zapain works quickly but only gives temporary relief. If the effect has worn off, be sure to wait at least four hours before taking the next dose.

Dosage

Each tablet or capsule of Zapain contains 500 mg paracetamol and 30 mg codeine. Zapain is available in packs of 100. Unless otherwise prescribed by your doctor, the general recommended dose is:

Adults: One to two tablets/capsules every four hours, with a maximum of eight tablets or capsules a day.
Young adults (aged 12-18): One to two tablets/capsules every six hours, with a maximum of eight tablets or capsules a day, for no more than three consecutive days. Zapain is not suitable for children under the age of 12.

Side effects

Zapain may cause side effects. Possible side effects include:

Constipation (this can be alleviated by drinking plenty of fluids and eating fibre-rich foods) Impaired reaction times: drowsiness, sleepiness, dizziness Stomach complaints (nausea, vomiting) See the package leaflet for a complete list of possible contraindications, warnings and other side effects.

Pregnancy/alcohol/driving ability

Zapain should only be used in consultation with a doctor if you are pregnant, are trying to conceive or are breastfeeding. The codeine in Zapain may affect your ability to react. Please be aware of this and do not participate in traffic if you are experiencing side effects. Alcohol and codeine amplify each other's effect, increasing the risk of drowsiness and dizziness. It is therefore advisable to stop or reduce alcohol consumption when you are taking Zapain.

Tags

[buy Zapain in Newport](#), [buy Zapain in Northern Ireland](#), [buy Zapain overnight](#), [order Zapain online in United Kingdom](#), [buy Zapain in Bradford](#), [buy Zapain in Scotland](#), [buy Zapain in Manchester](#), [Zapain online United Kingdom](#), [Zapain fast delivery](#), [Zapain online paypal](#), [Zapain tablets online in United Kingdom](#), [Zapain online pharmacy](#), [cheap Zapain online United Kingdom](#), [buy Zapain in Belfast](#), [buy Zapain online mastercard](#), [buy Zapain online in United Kingdom](#), [buy Zapain online with no prescription](#), [buy Zapain in Wales](#), [Zapain alternatives](#), [buy Zapain in Liverpool](#), [Zapain pills](#), [generic Zapain online](#), [buy Zapain online without prescription](#), [buy Zapain in Swansea](#), [cheapest Zapain online United Kingdom](#), [buy Zapain in England](#), [Zapain for sale](#),